

WHAT TO EXPECT AFTER TREATMENT

- For the majority of patients, downtime with **BELKYRA™** is minimal and occurs most commonly after their first treatment session.¹
- Apply ice or a cold pack to the treatment area for 10 to 15 minutes as needed.¹
- The most common side effects are swelling, bruising, pain, numbness, redness, areas of hardness, and tingling or itchiness in the treatment area. You can find detailed Important Safety Information on the reverse side.²
- While unlikely, talk to your doctor if you develop an asymmetric smile, facial muscle weakness, difficulty swallowing, or if symptoms worsen.²
- Remember, once your goals are achieved, retreatment with **BELKYRA™** is not expected.¹




PUT YOUR BEST CHIN FORWARD

Actual results with BELKYRA™.

Many people treated with **BELKYRA™** experienced visible improvement in the appearance of their chin profile in just 2 to 4 treatments.^{*2}

Up to 6 treatments may be administered.²

*Spaced at least one month apart. ¹Unretouched patient photos taken before and after 3 treatment sessions with **BELKYRA™**. Patient did not experience significant weight loss or gain between the start and end of treatment. Individual results will vary.¹



My Next Appointment

Date: _____ Time: _____

Before your next treatment, ask your doctor if there are any special instructions you should follow to prepare for your appointment.

IMPORTANT SAFETY INFORMATION ABOUT **BELKYRA™**



BELKYRA™ is an injectable prescription medicine used in adults to improve the appearance and profile of moderate to severe amounts of fat under the chin. **BELKYRA™** is only for use in a specific area under the chin and not for anywhere else on your body. **BELKYRA™** is only to be administered by a health professional who has been specially trained in the use of the product.

Do not use BELKYRA™ if you:

- are allergic to deoxycholic acid or any of the non-medicinal ingredients in the formulation.
- have an infection in your chin or neck area where the product will be injected.

To help avoid side effects and ensure proper use, talk to your health professional before BELKYRA™ is administered. Talk about any health conditions or problems you may have, including if you:

- have had, or plan to have, plastic surgery on your face, neck or chin, or if you have had other cosmetic treatments such as liposuction or neurotoxins (drugs sometimes used in the neck for cosmetic uses such as wrinkling or other medical reasons) in these areas.
- have, or have had, medical conditions in, on or near the neck.
- have difficulty swallowing.
- are pregnant or plan to become pregnant. It is not known if **BELKYRA™** can harm your unborn baby.
- are breastfeeding or plan to breastfeed as **BELKYRA™** may be found in breast milk.
- have any redness, swelling, pain or hard lumps in the area under the chin.
- have any medical condition (including high blood pressure, heart problems, coagulation (blood clotting) disorder or circulation problems, diabetes, lupus, have a compromised/weakened immune system, kidney problems, liver problems, or thyroid problems).
- have a history of skin darkening or lightening with medication.
- are taking corticosteroids used to treat joint pain and swelling.
- are younger than 18 years of age or over 65 years of age.

Tell your health professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

You may experience pain during the injection procedure. Your pain may last for several days after treatment. In the treatment area, you may also have bruising, swelling, numbness, redness, tingling or itchiness, and a sensation of warmth. You may experience a sensation of hardness across the treatment area or in small areas within the treatment area. Some patients reported headaches after receiving treatment.

BELKYRA™ can cause serious side effects, including: nerve injury (symptoms like uneven smile after treatment); trouble swallowing; low blood pressure immediately following treatment (symptoms like dizziness, fainting, lightheadedness); and high blood pressure (symptoms like headache, vision problems, irregular heartbeat).²

This is not a complete list of side effects. For any unexpected effects while taking **BELKYRA™**, contact your doctor or pharmacist.²

References: 1. Data on file, Allergan Inc. 2. [®]**BELKYRA™** Product Monograph. July 23, 2015.


belkyra™
deoxycholic acid injection

BELKYRA™ is a trademark of Kythera Biopharmaceuticals, Inc., an Allergan affiliate, used under license by Actavis Specialty Pharmaceuticals Co.
© 2016 Allergan. All rights reserved.

Allergan Aesthetics
an AbbVie company